

| LUNDI 9 DECEMBRE 2024 | | | |
|-----------------------|-----------------|---------------|--|
| ASCOCORYNE | CYLICNIUM | | (L.R. et C.Tulasne) Korf |
| BJERKANDERA | ADUSTA | | (Willdenow) Karsten |
| CALOCERA | VISCOSA | | (Pers.) Fr. |
| COPRINELLUS | MICACEUS | | (Bulliard) Vilgalys, Hopple et J.Johnson |
| COPRINOPSIS | LOTINAE | | (Picón) Picón |
| COPRINUS | COMATUS | | (O.F.Müller) Pers. |
| CORTINARIUS | COHABITANS | | Karsten |
| CORTINARIUS | SEMISANGUINEUS | | (Fr.) Gillet |
| CRATERELLUS | TUBAEFORMIS | | (Fr.) Quélet |
| CUPHOPHYLLUS | VIRGINEUS | | (Wulfen) Kovalenko |
| CYSTODERMA | AMIANTHINUM | | (Scopoli) Fayod |
| DALDINIA | CONCENTRICA | | (Bolton) Cesati et De Notaris |
| ENTOLOMA | CONFERENDUM | | (Britzelmayr) Noordeloos |
| ENTOLOMA | TENELLUM | | (J.Favre) Noordeloos |
| FUNALIA | GALLICA | | (Fr.) Bondartsev et Singer |
| GALERINA | HETEROCYSTIS | | (Atkinson) A.H.Smith et Singer |
| GALERINA | MARGINATA | | (Batsch) Kühner |
| GLOEOPHYLLUM | SEPIARIUM | | (Wulfen) Karsten |
| GYMNOPILUS | PENETRANS | | (Fr.) Murrill |
| HEBELOMA | HIEMALE | | Bresadola |
| HEBELOMA | MESOPHAEUM | | (Pers.) Quélet |
| HELVELLA | CRISPA | | (Scopoli) Fr. |
| HELVELLA | ELASTICA | | Bulliard |
| HELVELLA | LACUNOSA | | Afzelius |
| HYGROPHOROPSIS | AURANTIACA | | (Wulfen) Maire |
| HYPHOLOMA | FASCICULARE | | (Hudson) Kummer |
| INOCYBE | EUTHELES | | (Berkeley et Broome) Quélet |
| INOCYBE | GEOPHYLLA | | (Bulliard) Kummer |
| INOCYBE | GEOPHYLLA | var. LILACINA | (Peck) Gillet |
| INOCYBE | TENUICYSTIDIATA | | Horak et Stangl |
| LACCARIA | AFFINIS | | (Singer) Bon |
| LACCARIA | AMETHYSTINA | | M.C.Cooke |
| LACCARIA | BICOLOR | | (Maire) Orton |
| LACTARIUS | HEPATICUS | | Plowright |
| LENTINULA | EDODES | | (Berkeley) Pegler |
| LEPISTA | CAESPITOSA | | (Bresadola) Singer |
| LEPISTA | PERSONATA | | (Fr.) M.C.Cooke |
| LEPISTA | SORDIDA | | (C.F.Schumacher) Singer |
| MALVIPEZIA | PAULI | | (Medardi, Lantieri et Cacialli) Van Vooren |
| MELANOLEUCA | COGNATA | var. ROBUSTA | (J.E.Lange) Kühner |
| MELANOLEUCA | POLIOLEUCA | | (Fr.) Kühner et Maire |

| | | | |
|---------------|----------------|--|-----------------------------------|
| MERULIUS | TREMELLOSUS | | Schrader |
| MYCENA | CAPILLARIPES | | Peck |
| MYCENA | EPIPTERYGIA | | (Scopoli) Gray |
| MYCENA | INCLINATA | | (Fr.) Quélet |
| MYCENA | ROSEA | | (Bulliard) Gramberg |
| PANELLUS | STIPTICUS | | (Bulliard) Karsten |
| PAXILLUS | INVOLUTUS | | (Batsch) Fr. |
| PHOLIOTA | GUMMOSA | | (Lasch) Singer |
| PHOLIOTINA | DASYPUS | | (Romagnesi) P.-A. Moreau |
| PLEUROTUS | CORNUCOPIAE | | (Paulet) Quélet |
| PLUTEUS | CERVINUS | | (Schaeff.) Kummer |
| PSATHYRELLA | MULTIPEDATA | | (Peck) A.H. Smith |
| PSILOCYBE | CYANESCENS | | Wakefield |
| RHODOCOLLYBIA | BUTYRACEA | | (Bulliard) Lennox |
| RUSSULA | PUELLARIS | | Fr. |
| SCHIZOPHYLLUM | COMMUNE | | Fr. |
| STEREUM | INSIGNITUM | | Quélet |
| STROPHARIA | CAERULEA | | Kreisel |
| TEPHROCYBE | BOUDIERI | | (Kühner et Romagnesi) Derbsch |
| TOMENTELLA | LILACINOGRISEA | | Wakefield |
| TRAMETES | GIBBOSA | | (Pers.) Fr. |
| TRAMETES | OCHRACEA | | (Pers.) Gilbertson et Ryvarden |
| TRAMETES | VERSICOLOR | | (L.) Lloyd |
| TRICHOLOMA | SCALPTURATUM | | (Fr.) Quélet |
| TRICHOLOMA | SQUARRULOSUM | | Bresadola |
| TRICHOLOMA | TERREUM | | (Schaeff.) Kummer |
| VUILLEMINIA | COMEDENS | | (Nees) Maire |